

Partially dissipative hyperbolic systems, global existence in L^p spaces and relaxation limit¹

Timothée Crin-Barat

Chair of Computational Mathematics, University of Deusto, Spain

Tongji University, Shanghai - 25 April 2022



¹Results obtained in collaboration with Raphaël Danchin

1 Introduction

- General presentation
- Beauchard and Zuazua's method

2 Our approach

- New ideas
- Main results

3 Extensions

- Damped Baer-Nunziato system
- The HPC System
- Other

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

→ Model physical phenomena with finite speed of propagation or conservations laws, such as:

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

→ Model physical phenomena with finite speed of propagation or conservations laws, such as:

- The compressible Euler equation, Maxwell's equation, Einstein's equation, MHD equations, Yang-Mills equation, etc.

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

→ Model physical phenomena with finite speed of propagation or conservations laws, such as:

- The compressible Euler equation, Maxwell's equation, Einstein's equation, MHD equations, Yang-Mills equation, etc.
- Numerous applications: non-viscous fluid mechanics, kinetic theory, astrophysics, road traffic modelling, blood vessel circulation, etc.

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

→ Model physical phenomena with finite speed of propagation or conservations laws, such as:

- The compressible Euler equation, Maxwell's equation, Einstein's equation, MHD equations, Yang-Mills equation, etc.
- Numerous applications: non-viscous fluid mechanics, kinetic theory, astrophysics, road traffic modelling, blood vessel circulation, etc.

Question: When does the solution of (1) exists globally in time? And what happens when $\varepsilon \rightarrow 0$?

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

→ Model physical phenomena with finite speed of propagation or conservations laws, such as:

- The compressible Euler equation, Maxwell's equation, Einstein's equation, MHD equations, Yang-Mills equation, etc.
- Numerous applications: non-viscous fluid mechanics, kinetic theory, astrophysics, road traffic modelling, blood vessel circulation, etc.

Question: **When does the solution of (1) exists globally in time? And what happens when $\varepsilon \rightarrow 0$?**

More precisely we are interested in global strong and smooth solution being small perturbation of a constant state $\bar{V} \in \mathbb{R}^n$ such that $Q(\bar{V}) = 0$.

Introduction

We consider n -component quasilinear hyperbolic systems of the form:

$$\frac{\partial V}{\partial t} + \sum_{j=1}^d A^j(V) \frac{\partial V}{\partial x_j} = \frac{Q(V)}{\varepsilon}, \quad \text{where } V : \begin{cases} \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathcal{O}_V \subset \mathbb{R}^n \\ (t, x) \rightarrow V = V(t, x) \end{cases} \quad (1)$$

and $Q, A^j : \mathcal{O}_V \rightarrow \mathbb{R}^n$ are given n -vector valued smooth functions on \mathcal{O}_V .

→ Model physical phenomena with finite speed of propagation or conservation laws, such as:

- The compressible Euler equation, Maxwell's equation, Einstein's equation, MHD equations, Yang-Mills equation, etc.
- Numerous applications: non-viscous fluid mechanics, kinetic theory, astrophysics, road traffic modelling, blood vessel circulation, etc.

Question: **When does the solution of (1) exists globally in time? And what happens when $\varepsilon \rightarrow 0$?**

More precisely we are interested in global strong and smooth solution being small perturbation of a constant state $\bar{V} \in \mathbb{R}^n$ such that $Q(\bar{V}) = 0$.

→ There is a surprisingly strong connection between this question and problems related to control theory and Villani's hypocoercivity theory.

Dissipation term

For that issue, the 0-th order term Q , that act as a **dissipation** term, plays a crucial role.

Dissipation term

For that issue, the 0-th order term Q , that act as a **dissipation** term, plays a crucial role.

- In the case $Q \equiv 0$, local-in-time strong solutions may develop singularities (shock waves) in finite time (A. Majda, D. Serre).

Dissipation term

For that issue, the 0-th order term Q , that act as a **dissipation** term, plays a crucial role.

- In the case $Q \equiv 0$, local-in-time strong solutions may develop singularities (shock waves) in finite time (A. Majda, D. Serre).
- If $Q(V)$ acts directly on each component e.g. $Q(V) = -LV$ with $L > 0$
→ global existence + convergence to the equilibrium exponentially fast.
(T-T. Li)

Dissipation term

For that issue, the 0-th order term Q , that act as a **dissipation** term, plays a crucial role.

- In the case $Q \equiv 0$, local-in-time strong solutions may develop singularities (shock waves) in finite time (A. Majda, D. Serre).
- If $Q(V)$ acts directly on each component e.g. $Q(V) = -LV$ with $L > 0$
→ global existence + convergence to the equilibrium exponentially fast. (T-T. Li)
- In practice, a more reasonable assumption is that **the dissipation acts only on certain components of the system.**

Dissipation term

For that issue, the 0-th order term Q , that act as a **dissipation** term, plays a crucial role.

- In the case $Q \equiv 0$, local-in-time strong solutions may develop singularities (shock waves) in finite time (A. Majda, D. Serre).
- If $Q(V)$ acts directly on each component e.g. $Q(V) = -LV$ with $L > 0$
→ global existence + convergence to the equilibrium exponentially fast. (T-T. Li)
- In practice, a more reasonable assumption is that **the dissipation acts only on certain components of the system.**
- This can be written as:

$$Q(V) = \begin{pmatrix} 0_{\mathbb{R}^{n_1}} \\ q(V) \end{pmatrix} \text{ where } q(V) \in \mathbb{R}^{n_2}, n_1, n_2 \in \mathbb{N} \text{ and } n_1 + n_2 = n. \quad (2)$$

Dissipation term

For that issue, the 0-th order term Q , that act as a **dissipation** term, plays a crucial role.

- In the case $Q \equiv 0$, local-in-time strong solutions may develop singularities (shock waves) in finite time (A. Majda, D. Serre).
- If $Q(V)$ acts directly on each component e.g. $Q(V) = -LV$ with $L > 0$
→ global existence + convergence to the equilibrium exponentially fast. (T-T. Li)
- In practice, a more reasonable assumption is that **the dissipation acts only on certain components of the system**.
- This can be written as:

$$Q(V) = \begin{pmatrix} 0_{\mathbb{R}^{n_1}} \\ q(V) \end{pmatrix} \text{ where } q(V) \in \mathbb{R}^{n_2}, n_1, n_2 \in \mathbb{N} \text{ and } n_1 + n_2 = n. \quad (2)$$

- In the linear case, the system reads

$$\partial_t V + \sum_{j=1}^d A^j \partial_{x_j} V = \begin{pmatrix} 0 & 0 \\ 0 & -L \end{pmatrix} V \quad (3)$$

The question now is: how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?

The question now is: **how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?**

→ **Such a coercivity property is necessary to prove decay estimates and global existence**

The question now is: **how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?**

→ **Such a coercivity property is necessary to prove decay estimates and global existence**

As a toy-model, one may this damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + \nu v = 0. \end{cases} \quad (4)$$

The question now is: **how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?**

→ **Such a coercivity property is necessary to prove decay estimates and global existence**

As a toy-model, one may this damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + v = 0. \end{cases} \quad (4)$$

For this simple system, performing standard energy estimates leads to:

$$\frac{d}{dt} \|(u, v)\| + \|v\|_{L^2}^2 \leq 0 \rightarrow \text{no time-decay information on } u.$$

The question now is: **how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?**

→ **Such a coercivity property is necessary to prove decay estimates and global existence**

As a toy-model, one may this damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + v = 0. \end{cases} \quad (4)$$

For this simple system, performing standard energy estimates leads to:

$$\frac{d}{dt} \|(u, v)\| + \|v\|_{L^2}^2 \leq 0 \rightarrow \text{no time-decay information on } u.$$

Idea: consider the following perturbed functional

$$\mathcal{L}^2 = \|(u, v, \partial_x u, \partial_x v)\|_{L^2}^2 + \int_{\mathbb{R}} v \partial_x u,$$

which allows to recover dissipation properties on all the components.

The question now is: **how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?**

→ **Such a coercivity property is necessary to prove decay estimates and global existence**

As a toy-model, one may this damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + v = 0. \end{cases} \quad (4)$$

For this simple system, performing standard energy estimates leads to:

$$\frac{d}{dt} \|(u, v)\| + \|v\|_{L^2}^2 \leq 0 \rightarrow \text{no time-decay information on } u.$$

Idea: consider the following perturbed functional

$$\mathcal{L}^2 = \|(u, v, \partial_x u, \partial_x v)\|_{L^2}^2 + \int_{\mathbb{R}} v \partial_x u,$$

which allows to recover dissipation properties on all the components.

Indeed, after basic computations, we obtain

$$\frac{d}{dt} \mathcal{L}^2 + \|v\|_{L^2}^2 + \|(\partial_x u, \partial_x v)\|_{L^2}^2 \leq 0.$$

The question now is: **how can one recovers dissipative properties on all the components if the damping term is only present in some their equations?**

→ **Such a coercivity property is necessary to prove decay estimates and global existence**

As a toy-model, one may this damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + v = 0. \end{cases} \quad (4)$$

For this simple system, performing standard energy estimates leads to:

$$\frac{d}{dt} \|(u, v)\| + \|v\|_{L^2}^2 \leq 0 \rightarrow \text{no time-decay information on } u.$$

Idea: consider the following perturbed functional

$$\mathcal{L}^2 = \|(u, v, \partial_x u, \partial_x v)\|_{L^2}^2 + \int_{\mathbb{R}} v \partial_x u,$$

which allows to recover dissipation properties on all the components.

Indeed, after basic computations, we obtain

$$\frac{d}{dt} \mathcal{L}^2 + \|v\|_{L^2}^2 + \|(\partial_x u, \partial_x v)\|_{L^2}^2 \leq 0.$$

And since $\mathcal{L}^2 \sim \|(u, v, \partial_x u, \partial_x v)\|_{L^2}^2$, we can obtain time-decay estimates.

General case

General case

- Let's consider the following linear system:

$$\partial_t V + \sum_{j=1}^d A^j \partial_{x_j} V + LV = 0$$

where the A^j and L are constant symmetric matrices.

General case

- Let's consider the following linear system:

$$\partial_t V + \sum_{j=1}^d A^j \partial_{x_j} V + LV = 0$$

where the A^j and L are constant symmetric matrices.

- Applying the Fourier transform we get

$$\widehat{V}_t + iA(\xi)\widehat{V} + L\widehat{V} = 0 \quad \text{with} \quad A(\xi) \triangleq \sum_{j=1}^d A^j \xi_j.$$

General case

- Let's consider the following linear system:

$$\partial_t V + \sum_{j=1}^d A^j \partial_{x_j} V + LV = 0$$

where the A^j and L are constant symmetric matrices.

- Applying the Fourier transform we get

$$\widehat{V}_t + iA(\xi)\widehat{V} + L\widehat{V} = 0 \quad \text{with} \quad A(\xi) \triangleq \sum_{j=1}^d A^j \xi_j.$$

- In the 80's, Shizuta and Kawashima developed an algebraic criterion to ensure linear stability: the (SK) condition.

General case

- Let's consider the following linear system:

$$\partial_t V + \sum_{j=1}^d A^j \partial_{x_j} V + LV = 0$$

where the A^j and L are constant symmetric matrices.

- Applying the Fourier transform we get

$$\widehat{V}_t + iA(\xi)\widehat{V} + L\widehat{V} = 0 \quad \text{with} \quad A(\xi) \triangleq \sum_{j=1}^d A^j \xi_j.$$

- In the 80's, Shizuta and Kawashima developed an algebraic criterion to ensure linear stability: the (SK) condition.
- It is defined as follows:

Definition ((SK) condition)

$$\forall \xi \in \mathbb{R}^d, \quad \ker L \cap \{\text{eigenvectors of } A(\xi)\} = \{0\}. \quad (\text{SK})$$

General case

- Let's consider the following linear system:

$$\partial_t V + \sum_{j=1}^d A^j \partial_{x_j} V + LV = 0$$

where the A^j and L are constant symmetric matrices.

- Applying the Fourier transform we get

$$\widehat{V}_t + iA(\xi)\widehat{V} + L\widehat{V} = 0 \quad \text{with} \quad A(\xi) \triangleq \sum_{j=1}^d A^j \xi_j.$$

- In the 80's, Shizuta and Kawashima developed an algebraic criterion to ensure linear stability: the (SK) condition.
- It is defined as follows:

Definition ((SK) condition)

$$\forall \xi \in \mathbb{R}^d, \quad \ker L \cap \{\text{eigenvectors of } A(\xi)\} = \{0\}. \quad (\text{SK})$$

- Ensures that **the damping is strong enough** to prevent small solutions (perturbations of constant states) from developing singularities in finite time.

- Under this condition, they are able to prove the following decay estimates:

$$\begin{aligned}\|V^h(t)\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ce^{-\lambda t} \|V_0\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)}, \\ \|V^\ell(t)\|_{L^\infty(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ct^{-\frac{d}{2}} \|V_0\|_{L^1(\mathbb{R}^d, \mathbb{R}^n)}\end{aligned}\quad (5)$$

where V^h and V^ℓ correspond, respectively, to the high and low frequencies of the solution.

- Under this condition, they are able to prove the following decay estimates:

$$\begin{aligned}\|V^h(t)\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ce^{-\lambda t} \|V_0\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)}, \\ \|V^\ell(t)\|_{L^\infty(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ct^{-\frac{d}{2}} \|V_0\|_{L^1(\mathbb{R}^d, \mathbb{R}^n)}\end{aligned}\quad (5)$$

where V^h and V^ℓ correspond, respectively, to the high and low frequencies of the solution.

- Many results with this condition:

- Under this condition, they are able to prove the following decay estimates:

$$\begin{aligned}\|V^h(t)\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ce^{-\lambda t} \|V_0\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)}, \\ \|V^\ell(t)\|_{L^\infty(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ct^{-\frac{d}{2}} \|V_0\|_{L^1(\mathbb{R}^d, \mathbb{R}^n)}\end{aligned}\quad (5)$$

where V^h and V^ℓ correspond, respectively, to the high and low frequencies of the solution.

- Many results with this condition:
 - Hanouzet and Natalini '03, Ruggeri and Serre '04, **Yong '04**, Kawashima and Yong '04, '09 and **Kawashima and Xu '14**

- Under this condition, they are able to prove the following decay estimates:

$$\begin{aligned}\|V^h(t)\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ce^{-\lambda t} \|V_0\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)}, \\ \|V^\ell(t)\|_{L^\infty(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ct^{-\frac{d}{2}} \|V_0\|_{L^1(\mathbb{R}^d, \mathbb{R}^n)}\end{aligned}\quad (5)$$

where V^h and V^ℓ correspond, respectively, to the high and low frequencies of the solution.

- Many results with this condition:
 - Hanouzet and Natalini '03, Ruggeri and Serre '04, **Yong '04**, Kawashima and Yong '04, '09 and **Kawashima and Xu '14**
 - **Bianchini, Hanouzet and Natalini '07** and Kawashima and Xu '15.

- Under this condition, they are able to prove the following decay estimates:

$$\begin{aligned}\|V^h(t)\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ce^{-\lambda t} \|V_0\|_{L^2(\mathbb{R}^d, \mathbb{R}^n)}, \\ \|V^\ell(t)\|_{L^\infty(\mathbb{R}^d, \mathbb{R}^n)} &\leq Ct^{-\frac{d}{2}} \|V_0\|_{L^1(\mathbb{R}^d, \mathbb{R}^n)}\end{aligned}\quad (5)$$

where V^h and V^ℓ correspond, respectively, to the high and low frequencies of the solution.

- Many results with this condition:
 - Hanouzet and Natalini '03, Ruggeri and Serre '04, **Yong '04**, Kawashima and Yong '04, '09 and **Kawashima and Xu '14**
 - **Bianchini, Hanouzet and Natalini '07** and Kawashima and Xu '15.
- Here we develop a more explicit method partially based on Beauchard and Zuazua's paper (ARMA 2011).

Beauchard and Zuazua's method

Another point of view

Their paper (ARMA 2011) is based on the following classical proposition.

Proposition

Let A and L be two real matrices $n \times n$. The followings properties are equivalent:

- 1 the couple (A, L) satisfies the (SK) condition;*
- 2 the couple (A, L) satisfies the Kalman rank condition: the rank of $(L \quad AL \quad \dots \quad A^{n-1}L)$ is n ;*

Another point of view

Their paper (ARMA 2011) is based on the following classical proposition.

Proposition

Let A and L be two real matrices $n \times n$. The followings properties are equivalent:

- 1 the couple (A, L) satisfies the (SK) condition;*
- 2 the couple (A, L) satisfies the Kalman rank condition: the rank of $(L \quad AL \quad \dots \quad A^{n-1}L)$ is n ;*

→ link between the stability of these systems, control theory and Villani's hypocoercivity theory.

Inspired by this result, Beauchard and Zuazua designed the following Lyapunov functional:

$$\mathcal{L}^2 \triangleq \|V\|_{L^2}^2 + \int_{\mathbb{R}^d} \min(\rho, \rho^{-1}) \mathcal{I} \quad \text{where} \quad \mathcal{I} \triangleq \Im \sum_{k=1}^{n-1} \varepsilon_k (LA_\omega^{k-1} \widehat{V} \cdot LA_\omega^k \widehat{V})$$

for $n - 1$ positives parameters $\varepsilon_1, \dots, \varepsilon_{n-1}$.

Inspired by this result, Beauchard and Zuazua designed the following Lyapunov functional:

$$\mathcal{L}^2 \triangleq \|V\|_{L^2}^2 + \int_{\mathbb{R}^d} \min(\rho, \rho^{-1}) \mathcal{I} \quad \text{where} \quad \mathcal{I} \triangleq \Im \sum_{k=1}^{n-1} \varepsilon_k (LA_\omega^{k-1} \widehat{V} \cdot LA_\omega^k \widehat{V})$$

for $n - 1$ positives parameters $\varepsilon_1, \dots, \varepsilon_{n-1}$.

Differentiating in time \mathcal{L} and adjusting the coefficients ε_k , we obtain:

$$\frac{d}{dt} \mathcal{L}^2 + \mathcal{H} \leq 0 \quad \text{where} \quad \mathcal{H} \triangleq \frac{1}{2} \int_{\mathbb{R}^d} \sum_{k=0}^{n-1} \varepsilon_k \min(1, \rho^2) |LA_\omega^k \widehat{V}(\xi)|^2 d\xi.$$

Inspired by this result, Beauchard and Zuazua designed the following Lyapunov functional:

$$\mathcal{L}^2 \triangleq \|V\|_{L^2}^2 + \int_{\mathbb{R}^d} \min(\rho, \rho^{-1}) \mathcal{I} \quad \text{where} \quad \mathcal{I} \triangleq \Im \sum_{k=1}^{n-1} \varepsilon_k (LA_\omega^{k-1} \widehat{V} \cdot LA_\omega^k \widehat{V})$$

for $n - 1$ positives parameters $\varepsilon_1, \dots, \varepsilon_{n-1}$.

Differentiating in time \mathcal{L} and adjusting the coefficients ε_k , we obtain:

$$\frac{d}{dt} \mathcal{L}^2 + \mathcal{H} \leq 0 \quad \text{where} \quad \mathcal{H} \triangleq \frac{1}{2} \int_{\mathbb{R}^d} \sum_{k=0}^{n-1} \varepsilon_k \min(1, \rho^2) |LA_\omega^k \widehat{V}(\xi)|^2 d\xi.$$

- If $\varepsilon_1, \dots, \varepsilon_{n-1}$ are chosen small enough, then $\mathcal{L} \sim \|V\|_{L^2}^2$

Inspired by this result, Beauchard and Zuazua designed the following Lyapunov functional:

$$\mathcal{L}^2 \triangleq \|V\|_{L^2}^2 + \int_{\mathbb{R}^d} \min(\rho, \rho^{-1}) \mathcal{I} \quad \text{where} \quad \mathcal{I} \triangleq \Im \sum_{k=1}^{n-1} \varepsilon_k (LA_\omega^{k-1} \widehat{V} \cdot LA_\omega^k \widehat{V})$$

for $n - 1$ positives parameters $\varepsilon_1, \dots, \varepsilon_{n-1}$.

Differentiating in time \mathcal{L} and adjusting the coefficients ε_k , we obtain:

$$\frac{d}{dt} \mathcal{L}^2 + \mathcal{H} \leq 0 \quad \text{where} \quad \mathcal{H} \triangleq \frac{1}{2} \int_{\mathbb{R}^d} \sum_{k=0}^{n-1} \varepsilon_k \min(1, \rho^2) |LA_\omega^k \widehat{V}(\xi)|^2 d\xi.$$

- If $\varepsilon_1, \dots, \varepsilon_{n-1}$ are chosen small enough, then $\mathcal{L} \sim \|V\|_{L^2}^2$
- And, if the (SK) condition is satisfied then

$$\mathcal{H} \geq \kappa \min(1, |\xi|^2) \mathcal{L},$$

which brings us to

$$\frac{d}{dt} \mathcal{L} + \kappa \min(1, |\xi|^2) \mathcal{L} \leq 0$$

- At the linear level and under the condition (SK), this method allows to recover Kawashima's decay in a simple way.

- At the linear level and under the condition (SK), this method allows to recover Kawashima's decay in a simple way.
- More explicit method than Kawashima's \rightarrow very useful to study the high frequencies of non-linear systems.

- At the linear level and under the condition (SK), this method allows to recover Kawashima's decay in a simple way.
- More explicit method than Kawashima's \rightarrow very useful to study the high frequencies of non-linear systems.
- Their method covers some cases where condition (SK) is not verified.

- At the linear level and under the condition (SK), this method allows to recover Kawashima's decay in a simple way.
- More explicit method than Kawashima's \rightarrow very useful to study the high frequencies of non-linear systems.
- Their method covers some cases where condition (SK) is not verified.
- However, it does not tell the full story about the low frequencies, which turns out to be essential to study the relaxation limit issue.

Our approach

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces
 - *Hybrid* norm := norm with different regularity and/or Lebesgue indices in high and low frequencies.

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces
 - *Hybrid* norm := norm with different regularity and/or Lebesgue indices in high and low frequencies.
 - *Critical* := optimal conditions of smallness allowing to justify the global well-posedness of the system. → Corresponds to the largest space in which one can obtain the uniqueness of strong solutions.

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces
 - *Hybrid* norm := norm with different regularity and/or Lebesgue indices in high and low frequencies.
 - *Critical* := optimal conditions of smallness allowing to justify the global well-posedness of the system. → Corresponds to the largest space in which one can obtain the uniqueness of strong solutions.
- → More accurate information on the qualitative properties of the constructed solutions and a larger class of initial data under control.

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces
 - *Hybrid* norm := norm with different regularity and/or Lebesgue indices in high and low frequencies.
 - *Critical* := optimal conditions of smallness allowing to justify the global well-posedness of the system. → Corresponds to the largest space in which one can obtain the uniqueness of strong solutions.
- → More accurate information on the qualitative properties of the constructed solutions and a larger class of initial data under control.
- To have a better understanding of where our improvement fits in, we need to keep in mind the following embeddings:

$$H^s(s > \frac{d}{2} + 1) \hookrightarrow B_{2,1}^{\frac{d}{2}+1} \hookrightarrow \dot{B}_{p,1}^{\frac{d}{p}} \cap \dot{B}_{2,1}^{\frac{d}{2}+1} (p > 2) \hookrightarrow C_b^1.$$

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces
 - *Hybrid* norm := norm with different regularity and/or Lebesgue indices in high and low frequencies.
 - *Critical* := optimal conditions of smallness allowing to justify the global well-posedness of the system. → Corresponds to the largest space in which one can obtain the uniqueness of strong solutions.
- → More accurate information on the qualitative properties of the constructed solutions and a larger class of initial data under control.
- To have a better understanding of where our improvement fits in, we need to keep in mind the following embeddings:

$$H^s(s > \frac{d}{2} + 1) \hookrightarrow B_{2,1}^{\frac{d}{2}+1} \hookrightarrow \dot{B}_{p,1}^{\frac{d}{p}} \cap \dot{B}_{2,1}^{\frac{d}{2}+1} (p > 2) \hookrightarrow C_b^1.$$

- Since we work with homogeneous spaces, we can use their scaling properties and study the system with $\varepsilon = 1$. Then we recover the exact ε -dependency of the norm by scaling back.

Goal and ideas

- Work in the framework of *hybrid* and *critical* homogeneous Besov spaces
 - *Hybrid* norm := norm with different regularity and/or Lebesgue indices in high and low frequencies.
 - *Critical* := optimal conditions of smallness allowing to justify the global well-posedness of the system. → Corresponds to the largest space in which one can obtain the uniqueness of strong solutions.
- → More accurate information on the qualitative properties of the constructed solutions and a larger class of initial data under control.
- To have a better understanding of where our improvement fits in, we need to keep in mind the following embeddings:

$$H^s(s > \frac{d}{2} + 1) \hookrightarrow B_{2,1}^{\frac{d}{2}+1} \hookrightarrow \dot{B}_{p,1}^{\frac{d}{p}} \cap \dot{B}_{2,1}^{\frac{d}{2}+1} (p > 2) \hookrightarrow C_b^1.$$

- Since we work with homogeneous spaces, we can use their scaling properties and study the system with $\varepsilon = 1$. Then we recover the exact ε -dependency of the norm by scaling back.
- This allows to obtain optimal uniform estimates and therefore new results concerning the relaxation limit issue such as explicit convergence rates.

Littlewood-Paley decomposition

Littlewood-Paley decomposition

- We define $\dot{\Delta}_j$ as dyadic blocks such that $f \in \mathcal{S}'(\mathbb{R}^d)$

$$f = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j f \quad \text{and} \quad \text{supp}(\widehat{\dot{\Delta}_j f}) \subset \{\xi \in \mathbb{R}^d \text{ t.q. } \frac{3}{4}2^j \leq |\xi| \leq \frac{8}{3}2^j\}.$$

Littlewood-Paley decomposition

- We define $\dot{\Delta}_j$ as dyadic blocks such that $f \in \mathcal{S}'(\mathbb{R}^d)$

$$f = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j f \quad \text{and} \quad \text{supp}(\widehat{\dot{\Delta}_j f}) \subset \{\xi \in \mathbb{R}^d \text{ t.q. } \frac{3}{4}2^j \leq |\xi| \leq \frac{8}{3}2^j\}.$$

- The main motivation behind this decomposition is the following Bernstein inequality: $\forall k \in \mathbb{N}$

$$c2^{jk} \|\dot{\Delta}_j f\|_{L^p} \leq \|D^k \dot{\Delta}_j f\|_{L^p} \leq C2^{jk} \|\dot{\Delta}_j f\|_{L^p}.$$

Littlewood-Paley decomposition

- We define $\dot{\Delta}_j$ as dyadic blocks such that $f \in \mathcal{S}'(\mathbb{R}^d)$

$$f = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j f \quad \text{and} \quad \text{supp}(\widehat{\dot{\Delta}_j f}) \subset \{\xi \in \mathbb{R}^d \text{ t.q. } \frac{3}{4}2^j \leq |\xi| \leq \frac{8}{3}2^j\}.$$

- The main motivation behind this decomposition is the following Bernstein inequality: $\forall k \in \mathbb{N}$

$$C2^{jk} \|\dot{\Delta}_j f\|_{L^p} \leq \|D^k \dot{\Delta}_j f\|_{L^p} \leq C2^{jk} \|\dot{\Delta}_j f\|_{L^p}.$$

- The homogeneous Besov semi-norms are defined as follows:

$$\|f\|_{\dot{\mathbb{B}}_{p,1}^s} \triangleq \sum_{j \in \mathbb{Z}} 2^{js} \|\dot{\Delta}_j f\|_{L^p}.$$

Littlewood-Paley decomposition

- We define $\dot{\Delta}_j$ as dyadic blocks such that $f \in \mathcal{S}'(\mathbb{R}^d)$

$$f = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j f \quad \text{and} \quad \text{supp}(\widehat{\dot{\Delta}_j f}) \subset \{\xi \in \mathbb{R}^d \text{ t.q. } \frac{3}{4}2^j \leq |\xi| \leq \frac{8}{3}2^j\}.$$

- The main motivation behind this decomposition is the following Bernstein inequality: $\forall k \in \mathbb{N}$

$$C2^{jk} \|\dot{\Delta}_j f\|_{L^p} \leq \|D^k \dot{\Delta}_j f\|_{L^p} \leq C2^{jk} \|\dot{\Delta}_j f\|_{L^p}.$$

- The homogeneous Besov semi-norms are defined as follows:

$$\|f\|_{\dot{\mathbb{B}}_{p,1}^s} \triangleq \sum_{j \in \mathbb{Z}} 2^{js} \|\dot{\Delta}_j f\|_{L^p}.$$

- For a threshold $J_0 \in \mathbb{Z}$ and $s, s' \in \mathbb{R}$, we define:

Littlewood-Paley decomposition

- We define $\dot{\Delta}_j$ as dyadic blocks such that $f \in \mathcal{S}'(\mathbb{R}^d)$

$$f = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j f \quad \text{and} \quad \text{supp}(\widehat{\dot{\Delta}_j f}) \subset \{\xi \in \mathbb{R}^d \text{ t.q. } \frac{3}{4}2^j \leq |\xi| \leq \frac{8}{3}2^j\}.$$

- The main motivation behind this decomposition is the following Bernstein inequality: $\forall k \in \mathbb{N}$

$$C2^{jk} \|\dot{\Delta}_j f\|_{L^p} \leq \|D^k \dot{\Delta}_j f\|_{L^p} \leq C2^{jk} \|\dot{\Delta}_j f\|_{L^p}.$$

- The homogeneous Besov semi-norms are defined as follows:

$$\|f\|_{\dot{\mathbb{B}}_{p,1}^s} \triangleq \sum_{j \in \mathbb{Z}} 2^{js} \|\dot{\Delta}_j f\|_{L^p}.$$

- For a threshold $J_0 \in \mathbb{Z}$ and $s, s' \in \mathbb{R}$, we define:
- High-frequency norms: $\|f\|_{\dot{\mathbb{B}}_{2,1}^s}^h \triangleq \sum_{j \geq J_0} 2^{js} \|\dot{\Delta}_j f\|_{L^2},$

Littlewood-Paley decomposition

- We define $\dot{\Delta}_j$ as dyadic blocks such that $f \in \mathcal{S}'(\mathbb{R}^d)$

$$f = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j f \quad \text{and} \quad \text{supp}(\widehat{\dot{\Delta}_j f}) \subset \{\xi \in \mathbb{R}^d \text{ t.q. } \frac{3}{4}2^j \leq |\xi| \leq \frac{8}{3}2^j\}.$$

- The main motivation behind this decomposition is the following Bernstein inequality: $\forall k \in \mathbb{N}$

$$C2^{jk} \|\dot{\Delta}_j f\|_{L^p} \leq \|D^k \dot{\Delta}_j f\|_{L^p} \leq C2^{jk} \|\dot{\Delta}_j f\|_{L^p}.$$

- The homogeneous Besov semi-norms are defined as follows:

$$\|f\|_{\dot{\mathbb{B}}_{p,1}^s} \triangleq \sum_{j \in \mathbb{Z}} 2^{js} \|\dot{\Delta}_j f\|_{L^p}.$$

- For a threshold $J_0 \in \mathbb{Z}$ and $s, s' \in \mathbb{R}$, we define:

- High-frequency norms: $\|f\|_{\dot{\mathbb{B}}_{2,1}^s}^h \triangleq \sum_{j \geq J_0} 2^{js} \|\dot{\Delta}_j f\|_{L^2},$

- Low-frequency norms: $\|f\|_{\dot{\mathbb{B}}_{p,1}^{s'}}^\ell \triangleq \sum_{j \leq J_0} 2^{js'} \|\dot{\Delta}_j f\|_{L^p}.$

Crucial observations

Crucial observations

- In the previous approaches, the analysis of the low frequencies behaviour was not complete.

Crucial observations

- In the previous approaches, the analysis of the low frequencies behaviour was not complete.
- Back to the damped p -system (also the linearized compressible Euler system with damping):

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0. \end{cases} \quad (6)$$

A spectral analysis of the matrix

$$\begin{pmatrix} 0 & i\xi \\ i\xi & \frac{1}{\varepsilon} \end{pmatrix}.$$

shows that:

Crucial observations

- In the previous approaches, the analysis of the low frequencies behaviour was not complete.
- Back to the damped p -system (also the linearized compressible Euler system with damping):

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0. \end{cases} \quad (6)$$

A spectral analysis of the matrix

$$\begin{pmatrix} 0 & i\xi \\ i\xi & \frac{1}{\varepsilon} \end{pmatrix}.$$

shows that:

- the threshold between low and high frequencies is at $\frac{1}{2\varepsilon}$.

Crucial observations

- In the previous approaches, the analysis of the low frequencies behaviour was not complete.
- Back to the damped p -system (also the linearized compressible Euler system with damping):

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0. \end{cases} \quad (6)$$

A spectral analysis of the matrix

$$\begin{pmatrix} 0 & i\xi \\ i\xi & \frac{1}{\varepsilon} \end{pmatrix}.$$

shows that:

- the threshold between low and high frequencies is at $\frac{1}{2\varepsilon}$.
- In low frequencies (i.e. $|\xi| \ll \varepsilon^{-1}$), this matrix has two real eigenvalues asymptotically equal to $\frac{1}{\varepsilon}$ and $\varepsilon\xi^2$ for ξ close to 0;

Crucial observations

- In the previous approaches, the analysis of the low frequencies behaviour was not complete.
- Back to the damped p -system (also the linearized compressible Euler system with damping):

$$\begin{cases} \partial_t u + \partial_x v = 0, \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0. \end{cases} \quad (6)$$

A spectral analysis of the matrix

$$\begin{pmatrix} 0 & i\xi \\ i\xi & \frac{1}{\varepsilon} \end{pmatrix}.$$

shows that:

- the threshold between low and high frequencies is at $\frac{1}{2\varepsilon}$.
- In low frequencies (i.e. $|\xi| \ll \varepsilon^{-1}$), this matrix has two real eigenvalues asymptotically equal to $\frac{1}{\varepsilon}$ and $\varepsilon\xi^2$ for ξ close to 0;
- In high frequencies (i.e. $|\xi| \gg \varepsilon^{-1}$), two complex conjugate eigenvalues coexist, whose real parts are asymptotically equal to $\frac{1}{2\varepsilon}$.

This tells us that:

This tells us that:

- There should exist a damped mode in the low frequencies regime associated to the eigenvalue $\frac{1}{\varepsilon}$.

This tells us that:

- There should exist a damped mode in the low frequencies regime associated to the eigenvalue $\frac{1}{\varepsilon}$.
- As the eigenvalues are real in low frequencies, we expect to be able to work in L^p -space in this regime.

This tells us that:

- There should exist a damped mode in the low frequencies regime associated to the eigenvalue $\frac{1}{\varepsilon}$.
- As the eigenvalues are real in low frequencies, we expect to be able to work in L^p -space in this regime.
- The asymptotic behaviour of the solution when $\varepsilon \rightarrow 0$ is not so intuitive.

This tells us that:


- There should exist a damped mode in the low frequencies regime associated to the eigenvalue $\frac{1}{\varepsilon}$.
- As the eigenvalues are real in low frequencies, we expect to be able to work in L^p -space in this regime.
- The asymptotic behaviour of the solution when $\varepsilon \rightarrow 0$ is not so intuitive.
 - Naively, we expect that as the damping coefficient becomes larger the dissipation becomes more dominant.

This tells us that:

- There should exist a damped mode in the low frequencies regime associated to the eigenvalue $\frac{1}{\varepsilon}$.
- As the eigenvalues are real in low frequencies, we expect to be able to work in L^p -space in this regime.
- The asymptotic behaviour of the solution when $\varepsilon \rightarrow 0$ is not so intuitive.
 - Naively, we expect that as the damping coefficient becomes larger the dissipation becomes more dominant.
 - However, the so-called *overdamping* effect occurs: the decay rate behaves like $(\varepsilon, 1/\varepsilon)$.

This tells us that:

- There should exist a damped mode in the low frequencies regime associated to the eigenvalue $\frac{1}{\varepsilon}$.
- As the eigenvalues are real in low frequencies, we expect to be able to work in L^p -space in this regime.
- The asymptotic behaviour of the solution when $\varepsilon \rightarrow 0$ is not so intuitive.
 - Naively, we expect that as the damping coefficient becomes larger the dissipation becomes more dominant.
 - However, the so-called *overdamping* effect occurs: the decay rate behaves like $(\varepsilon, 1/\varepsilon)$.



- This phenomena makes things difficult when trying to study the relaxation limit of concrete systems.

→ To "overcome" this phenomenon, we choose a threshold between low and high frequencies that depends on the relaxation parameter ε .

→ To "overcome" this phenomenon, we choose a threshold between low and high frequencies that depends on the relaxation parameter ε .

- We define $J_\varepsilon = \lfloor -\log_2 \varepsilon \rfloor$ to have a better tracking of the relaxation parameter:

$$\|f\|_{\mathbb{B}_{2,1}^s}^{h,\varepsilon} \triangleq \sum_{j \geq J_\varepsilon} 2^{js} \|\dot{\Delta}_j f\|_{L^2} \quad \text{and} \quad \|f\|_{\mathbb{B}_{2,1}^s}^{\ell,\varepsilon} \triangleq \sum_{j \leq J_\varepsilon} 2^{js} \|\dot{\Delta}_j f\|_{L^2},$$

→ To "overcome" this phenomenon, we choose a threshold between low and high frequencies that depends on the relaxation parameter ε .

- We define $J_\varepsilon = \lfloor -\log_2 \varepsilon \rfloor$ to have a better tracking of the relaxation parameter:

$$\|f\|_{\mathbb{B}_{2,1}^s}^{h,\varepsilon} \triangleq \sum_{j \geq J_\varepsilon} 2^{js} \|\dot{\Delta}_j f\|_{L^2} \quad \text{and} \quad \|f\|_{\mathbb{B}_{2,1}^s}^{\ell,\varepsilon} \triangleq \sum_{j \leq J_\varepsilon} 2^{js} \|\dot{\Delta}_j f\|_{L^2},$$

- In the previous approaches, all the frequencies were "mixed" → the dissipative aspect in low frequencies was not fully exploited because the overall behaviour that emerges from a mixing of frequencies is always the "least good".

→ To "overcome" this phenomenon, we choose a threshold between low and high frequencies that depends on the relaxation parameter ε .

- We define $J_\varepsilon = \lfloor -\log_2 \varepsilon \rfloor$ to have a better tracking of the relaxation parameter:

$$\|f\|_{\mathbb{B}_{2,1}^s}^{h,\varepsilon} \triangleq \sum_{j \geq J_\varepsilon} 2^{js} \|\dot{\Delta}_j f\|_{L^2} \quad \text{and} \quad \|f\|_{\mathbb{B}_{2,1}^s}^{\ell,\varepsilon} \triangleq \sum_{j \leq J_\varepsilon} 2^{js} \|\dot{\Delta}_j f\|_{L^2},$$

- In the previous approaches, all the frequencies were "mixed" → the dissipative aspect in low frequencies was not fully exploited because the overall behaviour that emerges from a mixing of frequencies is always the "least good".
- Let us now transcribe this spectral analysis into concrete properties.

Low frequencies in a simple case

Back to the damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0 \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0, \end{cases}$$

Low frequencies in a simple case

Back to the damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0 \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0, \end{cases}$$

Defining the damped mode $w = v + \varepsilon \partial_x u$, the system can be rewritten

$$\begin{cases} \partial_t u - \varepsilon \partial_{xx}^2 u = -\partial_x w \\ \partial_t w + \frac{w}{\varepsilon} = -\varepsilon \partial_{xx}^2 v. \end{cases}$$

Low frequencies in a simple case

Back to the damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0 \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0, \end{cases}$$

Defining the damped mode $w = v + \varepsilon \partial_x u$, the system can be rewritten

$$\begin{cases} \partial_t u - \varepsilon \partial_{xx}^2 u = -\partial_x w \\ \partial_t w + \frac{w}{\varepsilon} = -\varepsilon \partial_{xx}^2 v. \end{cases}$$

→ This diagonalisation of the system directly exhibit the behaviours observed in the spectral analysis.

Low frequencies in a simple case

Back to the damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0 \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0, \end{cases}$$

Defining the damped mode $w = v + \varepsilon \partial_x u$, the system can be rewritten

$$\begin{cases} \partial_t u - \varepsilon \partial_{xx}^2 u = -\partial_x w \\ \partial_t w + \frac{w}{\varepsilon} = -\varepsilon \partial_{xx}^2 v. \end{cases}$$

→ This diagonalisation of the system directly exhibit the behaviours observed in the spectral analysis.

→ It is possible to study the two equations in a decoupled way as the source terms can be absorbed in the low-frequency regime:

$$\|\partial_x f\|_{B_{p,1}^s}^\ell \leq \|f\|_{B_{p,1}^s}^\ell$$

Low frequencies in a simple case

Back to the damped p-system:

$$\begin{cases} \partial_t u + \partial_x v = 0 \\ \partial_t v + \partial_x u + \frac{v}{\varepsilon} = 0, \end{cases}$$

Defining the damped mode $w = v + \varepsilon \partial_x u$, the system can be rewritten

$$\begin{cases} \partial_t u - \varepsilon \partial_{xx}^2 u = -\partial_x w \\ \partial_t w + \frac{w}{\varepsilon} = -\varepsilon \partial_{xx}^2 v. \end{cases}$$

→ This diagonalisation of the system directly exhibit the behaviours observed in the spectral analysis.

→ It is possible to study the two equations in a decoupled way as the source terms can be absorbed in the low-frequency regime:

$$\|\partial_x f\|_{B_{p,1}^s}^\ell \leq \|f\|_{B_{p,1}^s}^\ell$$

→ This allows to prove a well-posedness result with the low frequencies of the solution being bounded in L^p spaces, with $2 \leq p \leq 4$.

General case

General case

In the general case, the system can be rewritten as follows:

$$\begin{cases} \partial_t Z_1 + \sum_{k=1}^d \left(A_{1,1}^k(V) \partial_k Z_1 + A_{1,2}^k(V) \partial_k Z_2 \right) = 0, \\ \partial_t Z_2 + \sum_{k=1}^d \left(A_{2,1}^k(V) \partial_k Z_1 + A_{2,2}^k(V) \partial_k Z_2 \right) + \frac{L_2 Z_2}{\varepsilon} = 0. \end{cases}$$

General case

In the general case, the system can be rewritten as follows:

$$\begin{cases} \partial_t Z_1 + \sum_{k=1}^d \left(A_{1,1}^k(V) \partial_k Z_1 + A_{1,2}^k(V) \partial_k Z_2 \right) = 0, \\ \partial_t Z_2 + \sum_{k=1}^d \left(A_{2,1}^k(V) \partial_k Z_1 + A_{2,2}^k(V) \partial_k Z_2 \right) + \frac{L_2 Z_2}{\varepsilon} = 0. \end{cases}$$

We define the damped mode:

$$W \triangleq Z_2 + \varepsilon \sum_{k=1}^d L_2^{-1} \left(A_{2,1}^k(V) \partial_k Z_1 + A_{2,2}^k(V) \partial_k Z_2 \right) = -L_2^{-1} \partial_t Z_2.$$

General case

In the general case, the system can be rewritten as follows:

$$\begin{cases} \partial_t Z_1 + \sum_{k=1}^d \left(A_{1,1}^k(V) \partial_k Z_1 + A_{1,2}^k(V) \partial_k Z_2 \right) = 0, \\ \partial_t Z_2 + \sum_{k=1}^d \left(A_{2,1}^k(V) \partial_k Z_1 + A_{2,2}^k(V) \partial_k Z_2 \right) + \frac{L_2 Z_2}{\varepsilon} = 0. \end{cases}$$

We define the damped mode:

$$W \triangleq Z_2 + \varepsilon \sum_{k=1}^d L_2^{-1} \left(A_{2,1}^k(V) \partial_k Z_1 + A_{2,2}^k(V) \partial_k Z_2 \right) = -L_2^{-1} \partial_t Z_2.$$

The system can be rewritten

$$\begin{cases} \partial_t W + \frac{L_2 W}{\varepsilon} = g \\ \partial_t Z_1 - \varepsilon \sum_{k=1}^d \sum_{\ell=1}^d \bar{A}_{1,2}^k L_2^{-1} \bar{A}_{2,1}^\ell \partial_k \partial_\ell Z_1 = f \end{cases} \quad (7)$$

where f and g are controllable in the low-frequency regime.

General case

To study the equation of Z_1 , we have the following property

Lemma

Assume that $\forall k \in \{1, \dots, d\}$, $\bar{A}_{1,1}^k = 0$. The following assertions are equivalent:

- the system satisfy the (SK) condition at \bar{V} ;
- the operator $\mathcal{A} := \sum_{k=1}^d \sum_{\ell=1}^d \bar{A}_{1,2}^k L_2^{-1} \bar{A}_{2,1}^\ell \partial_k \partial_\ell$ is strongly elliptic.

General case

To study the equation of Z_1 , we have the following property

Lemma

Assume that $\forall k \in \{1, \dots, d\}$, $\bar{A}_{1,1}^k = 0$. The following assertions are equivalent:

- the system satisfy the (SK) condition at \bar{V} ;
- the operator $\mathcal{A} := \sum_{k=1}^d \sum_{\ell=1}^d \bar{A}_{1,2}^k L_2^{-1} \bar{A}_{2,1}^\ell \partial_k \partial_\ell$ is strongly elliptic.

→ We may study the equations of W and Z_1 separately, the former as a damped equation and the latter as a heat equation.

Main results

Recalling setting and assumptions

We look at multi-dimensional first order n -component systems in \mathbb{R}^d :

$$\frac{\partial V}{\partial t} + \sum_{k=1}^d A^k(V) \frac{\partial V}{\partial x_k} + \frac{LV}{\varepsilon} = 0,$$

such that:

- The maps A^k are symmetric valued \rightarrow hyperbolicity of the system.
- $L + {}^T L$ is nonnegative \rightarrow only *partial* dissipation occurs.
- An Hörmander's hypoellipticity-like condition is satisfied: the condition (SK): $\ker L \cap \{\text{eigenvectors of } A\} = \{0\}$

Main example of application:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho v) = 0, \\ \partial_t(\rho v) + \operatorname{div}(\rho v \otimes v) + \nabla P + \frac{1}{\varepsilon} \rho v = 0, \end{cases} \quad (8)$$

Well-posedness result

Theorem (Danchin, C-B '22)

Let $d \geq 1$, $p \in [2, 4]$ and $\varepsilon > 0$. There exists $k_p \in \mathbb{Z}$ and $c_0 = c_0(p) > 0$ such that for all $J_\varepsilon \triangleq \lfloor -\log_2 \varepsilon \rfloor + k_p$, if we assume

$$\|Z_0\|_{\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}}}^\ell + \varepsilon \|Z_0\|_{\dot{\mathbb{B}}_{2,1}^{\frac{d}{2}+1}}^h \leq c_0,$$

then the system admits a unique solution Z satisfying

$$X_{p,\varepsilon}(t) \lesssim \|Z_0\|_{\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}}}^\ell + \varepsilon \|Z_0\|_{\dot{\mathbb{B}}_{2,1}^{\frac{d}{2}+1}}^h \quad \text{for all } t \geq 0, \text{ and where}$$

$$\begin{aligned} X_{p,\varepsilon}(t) \triangleq & \varepsilon \|Z\|_{L_t^\infty(\dot{\mathbb{B}}_{2,1}^{\frac{d}{2}+1})}^h + \|Z\|_{L_t^1(\dot{\mathbb{B}}_{2,1}^{\frac{d}{2}+1})}^h + \varepsilon^{-\frac{1}{2}} \|Z_2\|_{L_t^2(\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}})} \\ & + \|Z\|_{L_t^\infty(\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}})}^\ell + \varepsilon \|Z_1\|_{L_t^1(\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}+2})}^\ell + \|Z_2\|_{L_t^1(\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}+1})}^\ell + \|W\|_{L_t^1(\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}})}. \end{aligned}$$

Decay estimates

Theorem (Danchin, C-B '22)

Assuming additionally that $Z_0 \in \dot{\mathbb{B}}_{2,\infty}^{-\sigma_1}$ for $\sigma_1 \in]-\frac{d}{2}, \frac{d}{2}]$ then there exists $C > 0$ such that

$$\|Z(t)\|_{\dot{\mathbb{B}}_{2,\infty}^{-\sigma_1}} \leq C \|Z_0\|_{\dot{\mathbb{B}}_{2,\infty}^{-\sigma_1}}, \quad \forall t \geq 0.$$

Moreover, if $\sigma_1 > 1 - d/2$,

$$\langle t \rangle \triangleq \sqrt{1+t^2}, \quad \alpha_1 \triangleq \frac{\sigma_1 + \frac{d}{2} - 1}{2} \quad \text{and} \quad C_0 \triangleq \|Z_0\|_{\dot{\mathbb{B}}_{2,\infty}^{-\sigma_1}}^\ell + \|Z_0\|_{\dot{\mathbb{B}}_{2,1}^{\frac{d}{2}+1}}^h,$$

then Z satisfies the following decay estimates:

$$\sup_{t \geq 0} \left\| \langle t \rangle^{\frac{\sigma+\sigma_1}{2}} Z(t) \right\|_{\dot{\mathbb{B}}_{2,1}^\sigma}^\ell \leq CC_0 \quad \text{if} \quad -\sigma_1 < \sigma \leq d/2 - 1,$$

$$\sup_{t \geq 0} \left\| \langle t \rangle^{\frac{\sigma+\sigma_1}{2} + \frac{1}{2}} Z_2(t) \right\|_{\dot{\mathbb{B}}_{2,1}^\sigma}^\ell \leq CC_0 \quad \text{if} \quad -\sigma_1 < \sigma \leq d/2 - 2,$$

$$\text{and} \quad \sup_{t \geq 0} \left\| \langle t \rangle^{2\alpha_1} Z(t) \right\|_{\dot{\mathbb{B}}_{2,1}^{\frac{d}{2}+1}}^h \leq CC_0.$$

Relaxation limit as $\varepsilon \rightarrow 0$

Relaxation limit as $\varepsilon \rightarrow 0$

As a direct consequence of the uniform bounds obtained in the existence theorem, we can treat the relaxation limit of the compressible Euler equations.

Relaxation limit as $\varepsilon \rightarrow 0$

As a direct consequence of the uniform bounds obtained in the existence theorem, we can treat the relaxation limit of the compressible Euler equations.

Introducing the slow-time variable $\tau = \varepsilon t$ and using the diffusive change of variable:

$$(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)(\tau, x) = \left(\rho, \frac{\rho u}{\varepsilon}\right)(t, x). \quad (9)$$

Relaxation limit as $\varepsilon \rightarrow 0$

As a direct consequence of the uniform bounds obtained in the existence theorem, we can treat the relaxation limit of the compressible Euler equations.

Introducing the slow-time variable $\tau = \varepsilon t$ and using the diffusive change of variable:

$$(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)(\tau, x) = \left(\rho, \frac{\rho u}{\varepsilon}\right)(t, x). \quad (9)$$

We get that $(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)$ satisfies:

$$\begin{cases} \partial_\tau \tilde{\rho}_\varepsilon + \nabla \cdot \tilde{J}_\varepsilon = 0, \\ \varepsilon^2 \left(\partial_\tau \tilde{J}_\varepsilon + \operatorname{div} \left(\frac{\tilde{J}_\varepsilon \otimes \tilde{J}_\varepsilon}{\tilde{\rho}_\varepsilon} \right) \right) + \nabla P(\tilde{\rho}_\varepsilon) + \tilde{J}_\varepsilon = 0. \end{cases} \quad (10)$$

Relaxation limit as $\varepsilon \rightarrow 0$

As a direct consequence of the uniform bounds obtained in the existence theorem, we can treat the relaxation limit of the compressible Euler equations.

Introducing the slow-time variable $\tau = \varepsilon t$ and using the diffusive change of variable:

$$(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)(\tau, x) = \left(\rho, \frac{\rho u}{\varepsilon}\right)(t, x). \quad (9)$$

We get that $(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)$ satisfies:

$$\begin{cases} \partial_\tau \tilde{\rho}_\varepsilon + \nabla \cdot \tilde{J}_\varepsilon = 0, \\ \varepsilon^2 \left(\partial_\tau \tilde{J}_\varepsilon + \operatorname{div} \left(\frac{\tilde{J}_\varepsilon \otimes \tilde{J}_\varepsilon}{\tilde{\rho}_\varepsilon} \right) \right) + \nabla P(\tilde{\rho}_\varepsilon) + \tilde{J}_\varepsilon = 0. \end{cases} \quad (10)$$

Then, formally, we have

$$-\tilde{J}_\varepsilon - \nabla P(\tilde{\rho}_\varepsilon) \xrightarrow{*} 0 \quad \text{et donc} \quad \partial_\tau \tilde{\rho}_\varepsilon - \Delta P(\tilde{\rho}_\varepsilon) \xrightarrow{*} 0.$$

Relaxation limit as $\varepsilon \rightarrow 0$

As a direct consequence of the uniform bounds obtained in the existence theorem, we can treat the relaxation limit of the compressible Euler equations.

Introducing the slow-time variable $\tau = \varepsilon t$ and using the diffusive change of variable:

$$(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)(\tau, x) = \left(\rho, \frac{\rho u}{\varepsilon}\right)(t, x). \quad (9)$$

We get that $(\tilde{\rho}_\varepsilon, \tilde{J}_\varepsilon)$ satisfies:

$$\begin{cases} \partial_\tau \tilde{\rho}_\varepsilon + \nabla \cdot \tilde{J}_\varepsilon = 0, \\ \varepsilon^2 \left(\partial_\tau \tilde{J}_\varepsilon + \operatorname{div} \left(\frac{\tilde{J}_\varepsilon \otimes \tilde{J}_\varepsilon}{\tilde{\rho}_\varepsilon} \right) \right) + \nabla P(\tilde{\rho}_\varepsilon) + \tilde{J}_\varepsilon = 0. \end{cases} \quad (10)$$

Then, formally, we have

$$-\tilde{J}_\varepsilon - \nabla P(\tilde{\rho}_\varepsilon) \xrightarrow{*} 0 \quad \text{et donc} \quad \partial_\tau \tilde{\rho}_\varepsilon - \Delta P(\tilde{\rho}_\varepsilon) \xrightarrow{*} 0.$$

And therefore, we expect $\tilde{\rho}_\varepsilon$ to converge to \mathcal{N} , the solution of the porous media equation:

$$\begin{cases} \partial_\tau \mathcal{N} - \Delta P(\mathcal{N}) = 0, \\ \mathcal{N}_{\tau=0} = \rho_0 \end{cases} \quad (11)$$

Relaxation result

Theorem (Danchin, C-B '21)

Let $d \geq 1$, $p \in [2, 4]$ and $\varepsilon > 0$. Let $\bar{\rho}$ be a strictly positive constant and $(\rho - \bar{\rho}, v)$ be the solution obtained with the previous theorem.

Let the positive function \mathcal{N}_0 such that $\mathcal{N}_0 - \bar{\rho}$ is small enough in $\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}}$, and let $\mathcal{N} \in \mathcal{C}_b(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}}) \cap L^1(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}+2})$ be the unique solution associated to the Cauchy problem:

$$\begin{cases} \partial_t \mathcal{N} - \Delta P(\mathcal{N}) = 0 \\ \mathcal{N}(0, x) = \mathcal{N}_0 \end{cases}$$

If we assume that

$$\|\tilde{\rho}_0^\varepsilon - \mathcal{N}_0\|_{\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}-1}} \leq C\varepsilon,$$

then

$$\|\tilde{\rho}^\varepsilon - \mathcal{N}\|_{L^\infty(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}-1})} + \|\tilde{\rho}^\varepsilon - \mathcal{N}\|_{L^1(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}+1})} + \left\| \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon} + \tilde{v}^\varepsilon \right\|_{L^1(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}})} \leq C\varepsilon.$$

Elements of proof

Elements of proof

The existence theorem can be applied to the Euler system under the Makino symmetrisation and assuming that $P(\rho) = A\rho^\gamma$ with $\gamma > 1$ and $A > 0$.

Elements of proof

The existence theorem can be applied to the Euler system under the Makino symmetrisation and assuming that $P(\rho) = A\rho^\gamma$ with $\gamma > 1$ and $A > 0$. Indeed, introducing the following 'sound speed':

$$c \triangleq \frac{2}{\gamma-1} \sqrt{\frac{\partial P}{\partial \rho}} = \frac{(4\gamma A)^{\frac{1}{2}}}{\gamma-1} \rho^{\frac{\gamma-1}{2}}. \quad (12)$$

Then, setting $\check{\gamma} = \frac{\gamma-1}{2}$ and $\tilde{c} = c - \bar{c}$, the Euler system reads

$$\begin{cases} \partial_t \tilde{c} + v \cdot \nabla \tilde{c} + \check{\gamma}(\tilde{c} + \bar{c}) \operatorname{div} v = 0, \\ \partial_t v + v \cdot \nabla v + \check{\gamma}(\tilde{c} + \bar{c}) \nabla \tilde{c} + \frac{1}{\varepsilon} v = 0. \end{cases} \quad (13)$$

Elements of proof

The existence theorem can be applied to the Euler system under the Makino symmetrisation and assuming that $P(\rho) = A\rho^\gamma$ with $\gamma > 1$ and $A > 0$. Indeed, introducing the following 'sound speed':

$$c \triangleq \frac{2}{\gamma-1} \sqrt{\frac{\partial P}{\partial \rho}} = \frac{(4\gamma A)^{\frac{1}{2}}}{\gamma-1} \rho^{\frac{\gamma-1}{2}}. \quad (12)$$

Then, setting $\check{\gamma} = \frac{\gamma-1}{2}$ and $\tilde{c} = c - \bar{c}$, the Euler system reads

$$\begin{cases} \partial_t \tilde{c} + v \cdot \nabla \tilde{c} + \check{\gamma}(\tilde{c} + \bar{c}) \operatorname{div} v = 0, \\ \partial_t v + v \cdot \nabla v + \check{\gamma}(\tilde{c} + \bar{c}) \nabla \tilde{c} + \frac{1}{\varepsilon} v = 0. \end{cases} \quad (13)$$

This system is symmetric and satisfies the (SK) condition.

Elements of proof

The existence theorem can be applied to the Euler system under the Makino symmetrisation and assuming that $P(\rho) = A\rho^\gamma$ with $\gamma > 1$ and $A > 0$. Indeed, introducing the following 'sound speed':

$$c \triangleq \frac{2}{\gamma-1} \sqrt{\frac{\partial P}{\partial \rho}} = \frac{(4\gamma A)^{\frac{1}{2}}}{\gamma-1} \rho^{\frac{\gamma-1}{2}}. \quad (12)$$

Then, setting $\check{\gamma} = \frac{\gamma-1}{2}$ and $\tilde{c} = c - \bar{c}$, the Euler system reads

$$\begin{cases} \partial_t \tilde{c} + v \cdot \nabla \tilde{c} + \check{\gamma}(\tilde{c} + \bar{c}) \operatorname{div} v = 0, \\ \partial_t v + v \cdot \nabla v + \check{\gamma}(\tilde{c} + \bar{c}) \nabla \tilde{c} + \frac{1}{\varepsilon} v = 0. \end{cases} \quad (13)$$

This system is symmetric and satisfies the (SK) condition. We use the diffusive change of variable:

$$(\tilde{c}^\varepsilon, \tilde{v}^\varepsilon)(\tau, x) = (c, \frac{v}{\varepsilon})(t, x).$$

Thus the pair $(\tilde{c}^\varepsilon, \tilde{v}^\varepsilon)$ satisfies

$$\begin{cases} \partial_t \tilde{c}^\varepsilon + \tilde{v}^\varepsilon \cdot \nabla \tilde{c}^\varepsilon + \check{\gamma} \tilde{c}^\varepsilon \operatorname{div} \tilde{v}^\varepsilon = 0, \\ \varepsilon^2 (\partial_t \tilde{v}^\varepsilon + \tilde{v}^\varepsilon \cdot \nabla \tilde{v}^\varepsilon) + \check{\gamma} \tilde{c}^\varepsilon \nabla \tilde{c}^\varepsilon + \tilde{v}^\varepsilon = 0. \end{cases} \quad (14)$$

Bounds for ρ

Bounds for ρ

Lemma

Let (\tilde{c}, v) be the solution from the existence theorem. Then, for all $t \geq 0$, $\tilde{\rho}^\varepsilon$ satisfies

$$\|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^\infty(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} + \|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^2(\mathbb{B}_{\rho,1}^{\frac{d}{p}+1})} + \frac{1}{\varepsilon} \left\| \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon} \right\|_{L_t^1(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} \leq c_0.$$

Bounds for ρ

Lemma

Let (\tilde{c}, v) be the solution from the existence theorem. Then, for all $t \geq 0$, $\tilde{\rho}^\varepsilon$ satisfies

$$\|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^\infty(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} + \|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^2(\mathbb{B}_{\rho,1}^{\frac{d}{p}+1})} + \frac{1}{\varepsilon} \left\| \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon} \right\|_{L_t^1(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} \leq c_0.$$

→ Key point: these bounds do not depend on the frequency regime → we do not have a problem when passing to the limit in our norms.

Bounds for ρ

Lemma

Let (\tilde{c}, v) be the solution from the existence theorem. Then, for all $t \geq 0$, $\tilde{\rho}^\varepsilon$ satisfies

$$\|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^\infty(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} + \|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^2(\mathbb{B}_{\rho,1}^{\frac{d}{p}+1})} + \frac{1}{\varepsilon} \left\| \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon} \right\|_{L_t^1(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} \leq c_0.$$

→ Key point: these bounds do not depend on the frequency regime → we do not have a problem when passing to the limit in our norms.

With this in hand we can estimate the difference of the solutions of:

$$\partial_t \mathcal{N} - \Delta P(\mathcal{N}) = 0 \quad \text{and} \quad \partial_t \tilde{\rho}^\varepsilon + \operatorname{div}(\tilde{\rho}^\varepsilon \tilde{v}^\varepsilon) = 0.$$

Bounds for ρ

Lemma

Let (\tilde{c}, v) be the solution from the existence theorem. Then, for all $t \geq 0$, $\tilde{\rho}^\varepsilon$ satisfies

$$\|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^\infty(\mathbb{B}_{\rho,1}^d)} + \|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^2(\mathbb{B}_{\rho,1}^{d+1})} + \frac{1}{\varepsilon} \left\| \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon} \right\|_{L_t^1(\mathbb{B}_{\rho,1}^d)} \leq c_0.$$

→ Key point: these bounds do not depend on the frequency regime → we do not have a problem when passing to the limit in our norms.

With this in hand we can estimate the difference of the solutions of:

$$\partial_t \mathcal{N} - \Delta P(\mathcal{N}) = 0 \quad \text{and} \quad \partial_t \tilde{\rho}^\varepsilon + \operatorname{div}(\tilde{\rho}^\varepsilon \tilde{v}^\varepsilon) = 0.$$

Using the damped mode $\tilde{W}^\varepsilon = \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\rho}$, the equation of $\tilde{\rho}^\varepsilon$ can be rewritten:

$$\partial_t \tilde{\rho}^\varepsilon - \Delta P(\tilde{\rho}^\varepsilon) = S^\varepsilon \quad \text{with} \quad S^\varepsilon = -\operatorname{div}(\tilde{\rho}^\varepsilon \tilde{W}^\varepsilon).$$

Bounds for ρ

Lemma

Let (\tilde{c}, v) be the solution from the existence theorem. Then, for all $t \geq 0$, $\tilde{\rho}^\varepsilon$ satisfies

$$\|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^\infty(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} + \|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^2(\mathbb{B}_{\rho,1}^{\frac{d}{p}+1})} + \frac{1}{\varepsilon} \left\| \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon} \right\|_{L_t^1(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} \leq c_0.$$

→ Key point: these bounds do not depend on the frequency regime → we do not have a problem when passing to the limit in our norms.

With this in hand we can estimate the difference of the solutions of:

$$\partial_t \mathcal{N} - \Delta P(\mathcal{N}) = 0 \quad \text{and} \quad \partial_t \tilde{\rho}^\varepsilon + \operatorname{div}(\tilde{\rho}^\varepsilon \tilde{v}^\varepsilon) = 0.$$

Using the damped mode $\tilde{W}^\varepsilon = \tilde{v}^\varepsilon + \frac{\nabla P(\tilde{\rho}^\varepsilon)}{\tilde{\rho}^\varepsilon}$, the equation of $\tilde{\rho}^\varepsilon$ can be rewritten:

$$\partial_t \tilde{\rho}^\varepsilon - \Delta P(\tilde{\rho}^\varepsilon) = S^\varepsilon \quad \text{with} \quad S^\varepsilon = -\operatorname{div}(\tilde{\rho}^\varepsilon \tilde{W}^\varepsilon).$$

The uniform bounds tell us that

$$\|\tilde{W}^\varepsilon\|_{L_t^1(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} = \mathcal{O}(\varepsilon) \quad \text{and} \quad \|\tilde{\rho}^\varepsilon - \bar{\rho}\|_{L_t^\infty(\mathbb{B}_{\rho,1}^{\frac{d}{p}})} = \mathcal{O}(1).$$

Therefore, using a standard product law and composition lemma, we obtain

$$\|S^\varepsilon\|_{L^1(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{d-1})} = \mathcal{O}(\varepsilon).$$

Therefore, using a standard product law and composition lemma, we obtain

$$\|S^\varepsilon\|_{L^1(\mathbb{R}^+; \dot{B}_{p,1}^{d-1})} = \mathcal{O}(\varepsilon).$$

Moreover, defining $\delta D^\varepsilon \triangleq \tilde{\rho}^\varepsilon - \mathcal{N}$, we have

$$\partial_t \delta D^\varepsilon - \Delta(P(\tilde{\rho}^\varepsilon) - P(\mathcal{N})) = S^\varepsilon.$$

Therefore, using a standard product law and composition lemma, we obtain

$$\|S^\varepsilon\|_{L^1(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}-1})} = \mathcal{O}(\varepsilon).$$

Moreover, defining $\delta D^\varepsilon \triangleq \tilde{\rho}^\varepsilon - \mathcal{N}$, we have

$$\partial_t \delta D^\varepsilon - \Delta(P(\tilde{\rho}^\varepsilon) - P(\mathcal{N})) = S^\varepsilon.$$

Using that:

- $\|\tilde{\rho}_0^\varepsilon - \mathcal{N}_0\|_{\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}-1}} \leq \varepsilon$

Therefore, using a standard product law and composition lemma, we obtain

$$\|S^\varepsilon\|_{L^1(\mathbb{R}^+; \dot{\mathbb{B}}_{p,1}^{\frac{d}{p}-1})} = \mathcal{O}(\varepsilon).$$

Moreover, defining $\delta D^\varepsilon \triangleq \tilde{\rho}^\varepsilon - \mathcal{N}$, we have

$$\partial_t \delta D^\varepsilon - \Delta(P(\tilde{\rho}^\varepsilon) - P(\mathcal{N})) = S^\varepsilon.$$

Using that:

- $\|\tilde{\rho}_0^\varepsilon - \mathcal{N}_0\|_{\dot{\mathbb{B}}_{p,1}^{\frac{d}{p}-1}} \leq \varepsilon$
- There exists a function H_1 vanishing at $\bar{\rho} = \bar{\mathcal{N}}$ such that

$$P(\tilde{\rho}^\varepsilon) - P(\bar{\rho}) = P'(\bar{\rho})(\tilde{\rho}^\varepsilon - \bar{\rho}) + H_1(\tilde{\rho}^\varepsilon)(\tilde{\rho}^\varepsilon - \bar{\rho}),$$

Therefore, using a standard product law and composition lemma, we obtain

$$\|S^\varepsilon\|_{L^1(\mathbb{R}^+; \mathbb{B}_{p,1}^{\frac{d}{p}-1})} = \mathcal{O}(\varepsilon).$$

Moreover, defining $\delta D^\varepsilon \triangleq \tilde{\rho}^\varepsilon - \mathcal{N}$, we have

$$\partial_t \delta D^\varepsilon - \Delta(P(\tilde{\rho}^\varepsilon) - P(\mathcal{N})) = S^\varepsilon.$$

Using that:

- $\|\tilde{\rho}_0^\varepsilon - \mathcal{N}_0\|_{\mathbb{B}_{p,1}^{\frac{d}{p}-1}} \leq \varepsilon$
- There exists a function H_1 vanishing at $\bar{\rho} = \bar{\mathcal{N}}$ such that

$$P(\tilde{\rho}^\varepsilon) - P(\bar{\rho}) = P'(\bar{\rho})(\tilde{\rho}^\varepsilon - \bar{\rho}) + H_1(\tilde{\rho}^\varepsilon)(\tilde{\rho}^\varepsilon - \bar{\rho}),$$

- The condition $P'(\bar{\rho}) > 0$, product law and composition lemma lead to

$$\|\delta D^\varepsilon\|_{L_T^\infty(\mathbb{B}_{p,1}^{\frac{d}{p}-1})} + \|\delta D^\varepsilon\|_{L_T^1(\mathbb{B}_{p,1}^{\frac{d}{p}+1})} \lesssim \varepsilon.$$

Extensions

Multifluid system

In a joint work with C. Burtea and J. Tan, we studied the following damped Baer-Nunziato system:

$$\left\{ \begin{array}{l} \partial_t \alpha_{\pm} + \mathbf{u} \cdot \nabla \alpha_{\pm} = \pm \frac{\alpha_+ \alpha_-}{2\mu + \lambda} (P_+(\rho_+) - P_-(\rho_-)), \\ \partial_t (\alpha_{\pm} \rho_{\pm}) + \operatorname{div} (\alpha_{\pm} \rho_{\pm} \mathbf{u}) = 0, \\ \partial_t (\rho \mathbf{u}) + \operatorname{div} (\rho \mathbf{u} \otimes \mathbf{u}) + \nabla P + \eta \rho \mathbf{u} = 0, \\ \rho = \alpha_+ \rho_+ + \alpha_- \rho_-, \\ P = \alpha_+ P_+(\rho_+) + \alpha_- P_-(\rho_-) \end{array} \right. \quad (BN)$$

Multifluid system

In a joint work with C. Burtea and J. Tan, we studied the following damped Baer-Nunziato system:

$$\left\{ \begin{array}{l} \partial_t \alpha_{\pm} + \mathbf{u} \cdot \nabla \alpha_{\pm} = \pm \frac{\alpha_+ \alpha_-}{2\mu + \lambda} (P_+(\rho_+) - P_-(\rho_-)), \\ \partial_t (\alpha_{\pm} \rho_{\pm}) + \operatorname{div} (\alpha_{\pm} \rho_{\pm} \mathbf{u}) = 0, \\ \partial_t (\rho \mathbf{u}) + \operatorname{div} (\rho \mathbf{u} \otimes \mathbf{u}) + \nabla P + \eta \rho \mathbf{u} = 0, \\ \rho = \alpha_+ \rho_+ + \alpha_- \rho_-, \\ P = \alpha_+ P_+(\rho_+) + \alpha_- P_-(\rho_-) \end{array} \right. \quad (BN)$$

With similar techniques as the one employed before, we are able to prove the global existence for this system as well as a result concerning the limit $\lambda, \mu \rightarrow 0$.

The HPC System

The HPC System

In a joint work with Q. He and L-Y. Shou, we studied the following hyperbolic-parabolic system:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla P(\rho) + \frac{1}{\varepsilon} \rho u - \mu \rho \nabla \phi = 0, \\ \partial_t \phi - \Delta \phi - a \rho + b \phi = 0, \end{cases} \quad x \in \mathbb{R}^d, \quad t > 0, \quad (\text{HPC})$$

The HPC System

In a joint work with Q. He and L-Y. Shou, we studied the following hyperbolic-parabolic system:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla P(\rho) + \frac{1}{\varepsilon} \rho u - \mu \rho \nabla \phi = 0, \\ \partial_t \phi - \Delta \phi - a \rho + b \phi = 0, \end{cases} \quad x \in \mathbb{R}^d, \quad t > 0, \quad (\text{HPC})$$

In this case, when $\varepsilon \rightarrow 0$, we show that the diffusive-rescaled solution of (HPC) converges strongly to the solution of the Keller-Segel system:

$$\begin{cases} \partial_t \rho - \operatorname{div}(\nabla P(\rho) - \mu \rho \nabla \phi) = 0, \\ \rho u = -\nabla P(\rho) + \mu \rho \nabla \phi, \\ -\Delta \phi - a \rho + b \phi = 0, \end{cases} \quad (\text{KS})$$

Other problem under investigation

- Jin-Xin system

Other problem under investigation

- Jin-Xin system
- Anisotropic systems e.g. 2D Boussinesq

Other problem under investigation

- Jin-Xin system
- Anisotropic systems e.g. 2D Boussinesq
- General system of any order (potentially anisotropic)

$$\partial_t V + A(D)V + L(D)V = 0, \quad \text{where}$$

- $A(D)$ is a skew-symmetric homogeneous Fourier multiplier of order α ,
- $L(D)$ is a partially elliptic homogeneous Fourier multiplier of order β .

Thank you for your attention! 谢谢!!